

Evaluation of Vitamin C and Some Selected Vitamins B in Human Milk and Infant Formulae for Different Storage Conditions

Klodiola DHAMO¹, Irida BREGOVINA², Lauresha SHABANI³

¹Pharmacy Department Aldent University Tirana Albania

²Biologist in Fier Albania

³Industrial Chemistry Department Faculty of Natural Science University of Tirana

Abstract:- Storage of human milk for further use in infant feeding, now is more frequently, as a result of the social- economic activities of breastfeeding mothers. Expressed breast milk is usually stored frozen for consumption at a later time. Currently, human milk is stored in containers with headspace, where the removal of oxygen should minimize the oxidative effects in compounds of stored breast milk during shelf life, and improve nutritional quality.

The purpose of this study was to analyze the quantity of some vitamins and compare any variation in vitamin C & some selected vitamins B (B1, B2, B3) in milk frozen in two different containers; one with air in the headspace versus vacuum sealed storage containers. Vitamin C content were analyzed using high-performance liquid chromatography (HPLC) Vitamin B1, B2, and B3 were also analyzed using HPLC. Fresh, mature, human milk donated by 4 women with healthy infants was used for analysis. Infant formula as powder and rehydrates liquid was also analyzed. According the results of analyses, the values of infant formula nutrient are generally higher than the human milk values.

Keywords: high-performance liquid chromatography, human milk, infant formula, Vitamin, B1, B2, B3, C.

I. INTRODUCTION

Breastfeeding women working outside the home commonly utilize a breast pumping device to provide milk for their infants. Human milk storage for use later in infant feeding is on the increase as a result of the economic activities of breastfeeding mothers. One reason for providing expressed milk to an infant is the anti-oxidant properties of human milk in the form of vitamins. It is known that fresh human milk has a higher antioxidant capacity than infant formula. [1] Expressed breast milk is usually stored frozen for consumption at a later time. It is known that changes occur in human milk stored either via refrigeration or frozen. Many nutrients, including vitamins, are susceptible to oxidation. Breast milk is subject to a strong peroxidation either at room temperature or at -20 degrees C. [2]. The oxidation of food products is well documented. Changes in product color, odor and taste are the most easily detected effects of oxidation in foods. [3]. Nutrient loss, rancidity, and microbial spoilage are all a part of milk degradation as well. This degradation, over time, is caused in part, by the presence of oxygen, and oxidation of the food components. [4]. Currently human milk is stored in containers with headspace. Removal of headspace oxygen should minimize the oxidative effects in stored breastmilk, prolong shelf life, and improve nutritional quality. The purpose of this study was to analyze and compare any differences in vitamin C and B1, B2, B3 in milk frozen in containers with ambient air in the headspace versus vacuum sealed storage containers. [5].

II. EXPERIMENTAL

Fresh, mature, human milk donated by four women with healthy infants was used for analysis. Infant formula from powder and liquid was also analyzed. Baseline analysis of each sample was completed within 24 hours of collection for Vitamin C, B1, B2, and B3. Each sample was divided into two sets, set S (standard) and V (vacuum) with aliquots of 1 ounces each: i.e., Day 10, Day 20, and Day 40. A 20 ml sample was removed from each aliquot and tested for the nutrients of interest. Half the samples were then vacuum sealed and placed in the freezer at -20 °C. The other half of the samples were sealed with ambient air and placed in freezer at -20 °C. After 10 days, one aliquot from each group was removed for analysis. At 20 days and 40 days the process was repeated.

Vitamin C content was analyzed using [6]. The samples were protected from light by wrapping tubes and flasks with aluminum foil and preparing the samples in a darkened room. Three hundred µl of milk mixed with 300ul of 0.56% meta-phosphoric acid solution were added to the same centrifuge and filtration tube, which was shaken for 30 seconds and centrifuged at 10°C (10 minutes, 3000 X g). Ascorbic acid was identified by comparing the retention time of the sample peak with that of the ascorbic standard at 254 nm. Quantification

was carried out using external standardization. Vitamin B1, B2, and B3 were analyzed using HPLC as described by [7] The samples were protected from light by wrapping tubes and flasks with aluminum foil and preparing the samples in a darkened room. 10.5 grams of sample were weighed into a 50 ml centrifuge tube. 1 g of TCA solid and a magnetic stirring bar were added. The mixture was stirred for 10 min over a magnetic stirring plate, then centrifuged for 10 minutes at 1250 g to separate the two phases. Three ml of 4% TCA was added to the solid residue obtained and mixed for 10 minutes and then centrifuged. The solid-phase was discarded. The two acid extracts were combined in a 10 ml volumetric flask and the volume was filled with 4% TCA. Acid extracts were filtered through a .45um filter and injected into the HPLC system. This process was repeated for fresh milk stored under refrigeration for 48 hours, with measurement of nutrients occurring at 12, 24, and 48 hour intervals. Descriptive statistics were calculated and nutrient mean values were compared using paired t-tests. Statistical significance was determined when $p < .01$.

III. RESULTS AND DISCUSSIONS

Descriptive statistics were calculated and nutrient mean values were compared using paired t-tests. Statistical significance was determined when $p < .01$.

The baseline values for each sample and nutrient are listed in TABLE 1. These values are highly variable, as would be expected in human milk. The infant formula nutrient values are generally higher than the human milk values. The Vitamin C levels ranged from 32 mg/ml to a high of 89 mg/ml. The range for vitamin B1 was 0.032 mg/ml to 0.654 mg/ml. The range for vitamin B2 was 0.156 mg/ml to 1.015 mg/ml. The range for vitamin B3 was 1.25 mg/ml to 7.10 mg/ml

Table 1. Baseline values by nutrient and sample

	C (mg/L)	B1 (mg/L)	B2 (mg/L)	B3 (mg/L)
HM1	51	0.082	0.511	1.25
HM2	87	0.185	0.156	5.33
HM3	32	0.032	0.456	4.28
HM4	89	0.096	0.396	1.94
IF1	61	0.476	0.924	7.10
IF2	81	0.654	0.743	6.76
Average	66.8	0.254	0.531	4.443

Due to the high variability of the values for human milk the mean values were used for comparison. The decrease in levels of vitamin A and E was higher in the S group than in the V group and was statistically significantly different.

The baseline values and the final values for each group are listed in TABLE 2.

The mean values for the human milk samples are depicted in TABLE 3.

Table 2. Values at baseline and 40 days.

	C(mg/L)			B1(mg/L)			B2(mg/L)			B3(mg/L)		
	Baseline	S40	V40	Baseline	S40	V40	Baseline	S40	V40	Baseline	S40	V40
HM1	453	337	441	0.082	0.059	0.082	0.511	0.204	0.236	1.25	0.54	0.95
HM2	694	602	634	0.185	0.119	0.147	0.156	0.065	0.185	5.33	4.09	4.19
HM3	196	179	182	0.032	0.000	0.000	0.456	0.389	0.401	4.28	3.68	4.01
HM4	302	267	297	0.096	0.048	0.068	0.396	0.111	0.109	1.94	1.41	1.52
IF1	712	482	484	0.476	0.476	0.496	0.924	0.821	0.901	7.10	6.489	6.564
IF2	1147	1038	1038	0.654	0.359	0.486	0.743	0.699	0.703	6.76	5.680	5.897
Average	584	484	512	0.254	0.176	0.213	0.531	0.381	0.422	4.443	3.648	3.855

Table 3. Mean values of vitamins in human milk.

Vitamin	Baseline	10 days		20 days		40 days	
		S Group	V Group	S Group	V Group	S Group	V Group
C mg/ml	0.058	0.041	0.052	0.034	0.042	0.018	0.041
B1 mcg/ml	0.096	0.073*	0.088	0.064*	0.080	0.045*	0.067*
B2 mcg/ml	0.315	0.281**	0.300	0.255**	0.286	0.219**	0.261**#
B3 mcg/ml	0.283	0.249*	0.267	0.223*	0.246	0.195*	0.229

*Statistically significantly below baseline at $p < 0.01$.

**Statistically significantly below baseline at $p < 0.001$.

IV. CONCLUSION

All of the nutrient levels dropped for the duration of the time in storage in both groups but consistently less in the V group. For example, at baseline, most subjects had adequate levels of vitamins at baseline to meet the RDI for their infant. However, by the end of the 40 days storage time not all samples contained the RDI. by the end of the 40 days storage time not all samples contained the RDI for vitamin C of 50 mg/day for infants of less than 1 year of age. In particular, vitamin C content from the S group was reduced to non detectable levels in four of the human milk samples by the last measurement, while the V group all had Nutrients were better preserved by vacuum sealing, providing nutrient levels near the RDI for infants under the age of 1 year. Women around the world are storing their milk frozen for times of separation from their infants. [8]. Providing a storage system that preserves the nutrients in frozen human milk is critical for the long-term health of those infants fed with frozen milk.

REFERENCES

- [1]. JK. FRIEL, SM. MARTIN, M. LANGDON, GR. HERZBERG GR BUETTNER.. Milk from mothers of both premature and full-term infants provides better antioxidant protection than does infant formula. *Pediatric Research*. 51(5): 2002,612-618.
- [2]. D. TUROLI, G. TESTOLIN, R. ZANINI, R. BELLU Determination of oxidative status in breast and formula milk. *Acta Paediatr.*. 90(7): 2001,813-5.
- [3]. DA. CLARE, WS. BANG, G. CARTWRIGHT, MA.DRAKE, P. CORONEL, J. SIMUNOVIC Comparison of sensory, microbiologic, and biochemical parameters of microwave versus indirect UHT fluid skim milk during storage. *J Dairy Science.*. 88: 2005,4172- 4182.
- [4]. LJ. WHITED, BH. HAMMOND, KW. CHAPMAN, KJ. BOOR Vitamin A degradation and light-oxidized flavor defects in milk. *J Dairy Science.*. 85: 2002,351-354.
- [5]. JL. CHAVEZ-SERVIN, AI. CASTELLOTE, MC. LOPEZ-SABATER. Simultaneous analysis of Vitamins A and E in infant milk-based formulae by normal-phase highperformance liquid chromatography-diode array detection using a short narrowbore column. *J of Chromatography*. 1122: 2006,138.143.
- [6]. M. ROMEU-NADAL, S. MORERA-PONS, AI. CASTELLOTE, MC. LOPEZ-SABATER Rapid highperformance liquid chromatographic method for Vitamin C determination in human milk versus an enzymatic method. *J Chromatography.*. 830: 2006 ,41-46.
- [7]. S. ALBALA-HURTADO, MT VECIANNA-NOGUES, M. IZQUIERDO-PULIDO, A. MARINE-FONT, Determination of water-soluble vitamins in infant milk by high-performance liquid chromatography. 778: 1997,247-253.
- [8]. EL. EL DIN ZM, SA. GHAFAR, EK. EL GABRY, WA. FAHMI, RF. BEDAIR. Is stored expressed breast milk an alternative for working Egyptian mothers? *Eastern Mediterranean Health J*. 10(6): 2004, 815-821.